CONTENTS

Thermodynamic parameters of ionization of Kojic acid and its coordinating behaviour towards transition metals
S.A. Sallam, S.S. Haggag and M.S. Masoud (Alexandria, Egypt) 1
Thermal behaviour of high α-cellulose pulp from fast growing plant species and its acetylation product
C.N. Saikia, N.N. Dutta and J.N. Baruah (Jorhat, India) and N.N. Dass (Dibrugarh, India) . . 9
The electrical conductivity of benzaldazine and its *o*-, *m*- and *p*-hydroxy substituents
M.A. Ahmed (Giza, Egypt), A. El-Sayed Badr, M. El-Desoky Moustafa and M. Sha'aban Salah (Cairo, Egypt) 25
Prediction of excess functions of some multicomponent organic mixtures of hydrocarbons with a group contribution model
J.A. Gonzalez, I. Garcia, J.C. Cobos and C. Casanova (Valladolid, Spain) 31
Thermal, spectral, ESR and magnetic studies of some copper(II) thiosemicarbazone complexes
S. Laly and G. Parameswaran (Kerala, India) 43
Kinetic study by mass spectrometry of the decomposition of basic cobalt carbonate
M.M. Romero (Holguin, Cuba), J.C. Llópiz (Habana, Cuba), A. Jerez, C. Pico and M.L. Veiga (Madrid, Spain) 53
Reaction entropy for the electroreduction of 2-phenyl-3-arylimino-3*H*-indoles in various aprotic solvents
G. Marrosu, F. Rodante and A. Trazza (Roma, Italy) and L. Greci (Ancona, Italy) 59
Some problems concerning the temperature integral in non-isothermal kinetics
Part I. Generalities and some simple applications
E. Urbanovici (Județul Covasna, Romania) and E. Segal (Bucharest, Romania) 71
Translational heat capacity in small cubic boxes: a possible contribution to temperature heat capacity maximum in Pd-hydrogen systems
Z. Slanina (Mainz, F.R.G) 89
Excess volumes for binary mixtures formed by 2,5,8,11-tetraoxadodecane or 2,5,8,11,14-pentaoxapentadecane with homologous *n*-alkanes
T. Treszczanowicz, G.C. Benson and B.C-Y. Lu (Ottawa, Ont., Canada) 95
A detailed study of the condensation of the $Ca(H_2PO_4)_2 \cdot H_2O$–$CaHPO_4 \cdot 2H_2O$ system under thermal treatment
T.C. Vaimakis, P.J. Pomonis and A.T. Sdoukos (Ioannina, Greece) 103
Kinetics of *n*-heptane desorption from pelletized zeolite 5A
A Palermo and D.G. Löffler (Mar del Plata, Argentina) 115
Enthalpies of mixing at 298.15 K of a methyl alkanoate (from acetate to pentanoate) with *n*-alkanes (*n*-tridecane and *n*-pentadecane)
J. Ortega, J.S. Matos and J.A. Peña (Canary Islands, Spain) 121
Analysis of excess molar volumes of binary mixtures of toluene with the hexanol isomers at 298.15 K
J.S. Matos and J. Ortega (Canary Islands, Spain) and M.V. Garcia (Madrid, Spain) 127
A thermoanalytical study of the synthesis and oxidation of $NaCrO_2$
C.P.J. van Vuuren (Pretoria, South Africa) 135
Aqueous micellar solutions of some *N*-alkylnicotinamide and *N*-alkylpyridinium halide surfactants: apparent and partial molar enthalpies
E. Fisicaro, E. Pelizzetti and M. Barbieri (Parma, Italy), P. Savarino and G. Viscardi (Torino, Italy) 143
Calculation of site enthalpy for binding of ligands and protons to macromolecules
E. Fisicaro and A. Braibanti (Parma, Italy), J.D. Lamb and J.L. Oscarson (Provo, UT, U.S.A.) 161
Group additivity scheme for calculating the chemical thermodynamic properties of gaseous polycyclic aromatic hydrocarbons containing five-membered rings
N.F. Moiseeva and O.V. Dorofeeva (Moscow, U.S.S.R.) 179

Thermal and spectral study of 5-chloro-4-β-glycopyranosylaminopyrimidine derivatives
A. Sánchez, M. Nogueras, M^a.L. Quijano, M. Melguizo and R. López (Jaén, Spain) 187
Thermal decomposition of 4,6-diamino-1,2-dihydro-2-thiopyrimidine and its complexes with Co^{II}, Co^{III}, Ni^{II}, and Cu^{II} ions
M.D. Gutierrez, M.A. Romero, R. López, M. Nogueras and A. Sánchez (Jaén, Spain) 197
Thermal expansion studies on some rare-earth pyrohafnates by high temperature X-ray powder diffractometry
K.V.G. Kutty, S. Rajagopalan and R. Asuvathraman (Kalpakkam, India) 205
Thermal decomposition of tetrazole-containing polymers. I. Poly-5-vinyltetrazole thermolysis
S.V. Levchik, E.E. Bolvanovich, A.I. Lesnikovich, O.A. Ivashkevich, P.N. Gaponik and S.V. Vyazovkin (Minsk, U.S.S.R.) 211
Thermal decomposition of hydrated selenites of trivalent lanthanides and yttrium
M. Ionashiro, C.B. Melios, C.A. Ribeiro, M. Spirandeli Crespi and I. Giolito (São Paulo, Brazil) 223
Thermodynamics of associated solutions. Excess enthalpies for acetic acid–non-polar component mixtures
I. Nagata (Ishikawa, Japan) 233
Thermodynamics of associated solutions. Vapour–liquid equilibrium and excess enthalpy for acetic acid–polar unassociated component mixtures
I. Nagata and T. Tanimura (Ishikawa, Japan) 241
The properties of salt-filled sodalites. I. Thermal decomposition reactions of hydroxoborate sodalite
J.Ch. Buhl (Münster, F.R.G.) and S. Luger (Lahn, F.R.G.) 253

Review
Application of thermal analysis for investigating liquid fuels, petroleum- and coke-chemical products
D.D. Rustschev (Sofia, Bulgaria) 261

Notes
Studies on double selenates. IX. Thermal decomposition of lanthanide of yttrium and magnesium double selenates
P.R. de Ávila Agostini and E. de Castilho Agostini (Alfenas, Brazil), I. Giolito (São Paulo, Brazil) and M. Ionashiro (Araraquara, Brazil) 273
Studies on double selenates. XI. Thermal decomposition of lanthanides of the ceric group and lithium double selenates
P.M. Cruz, M.S. Crespi, C.A. Ribeiro and M. Ionashiro (Araraquara, Brazil) and I. Giolito (São Paulo, Brazil) 283
Estimation of the degree of hydration in modified ordinary Portland cement pastes by differential scanning calorimetry. Part II
B.E.I. Abdelrazig, D.G. Bonner and D.V. Nowell (Hatfield, Great Britain), P.J. Egan and J.M. Dransfield (Birmingham, Great Britain) 291

Author Index 297

Subject Index 299